


(Free and download) Morphology and histology of plants, designed especially as a guide to plant-analysis and classification, and as an introduction to pharmacognosy and vegetable physiology

## **Morphology and histology of plants, designed especially as a guide to plant-analysis and classification, and as an introduction to pharmacognosy and vegetable physiology**

*Henry Hurd Rusby*

*ePub | \*DOC | audiobook | ebooks | Download PDF*



 [Download](#)

 [Read Online](#)

1899-01-01 8.50 x 1.03 x 5.50l, #File Name: B003YJGG3E412 pages | File size: 31.Mb

**Henry Hurd Rusby : Morphology and histology of plants, designed especially as a guide to plant-analysis and classification, and as an introduction to pharmacognosy and vegetable physiology** before purchasing it in order to gage whether or not it would be worth my time, and all praised Morphology and histology of plants, designed

especially as a guide to plant-analysis and classification, and as an introduction to pharmacognosy and vegetable physiology:

This historic book may have numerous typos and missing text. Purchasers can download a free scanned copy of the original book (without typos) from the publisher. Not indexed. Not illustrated. 1899 Excerpt: ...an indumentum in the form of granular or detached scaly masses. When the matter of such masses is more thinly distributed, appearing in the form of a powder rather than a scurf, the surface is called Pulverulent. A Pubescent surface is a hairy surface which is not readily distinguished as pertaining to any one of the other specific classes. If the hairs of a pubescent surface are very short and fine, so that the consequent roughness is reduced to a minimum, the surface is called Puberulent. If a hairy indumentum is fine and of an ashy-gray color, the hairs not arranged in any regular direction, the surface is called Cinereous. If the hairs all lie in one direction, are closely appressed and have a shiny or silky lustre, the surface is called Sericeous. If this lustre is intensified and of a strongly whitish color, whether the trichomes be hairs or scales, the surface is denominated Argenteous. Such hairs as are capable of producing a sericeous surface are themselves denominated sericeous or silky, even though they be in insufficient numbers to impart this character to the general surface. A surface tending toward the sericeous, but not sufficiently pronounced, is called Canescent. When there is a dense covering of more or less elongated and matted hairs, the surface is called Tomentose. When such a covering is thin, its hairs less elongated, it is called Tomentellate. When there is a covering of thinly distributed, elongated, moderately soft hairs, which are not closely appressed, the surface is Pilose. When hairs are similarly distributed, but are elongated and coarse, the surface is Hirsute. When similar coarse hairs are rather stiff, lie in one direction, somewhat appressed, and particularly when each develops from an elevated base, the surface is Strigose. A s...